

DIRECTIONS FOR USE

THIS INSECT CONTAINS COMPLETE DIRECTIONS FOR USE. PLEASE READ CAREFULLY AND DO NOT DISCARD UNTIL ALL PRODUCTS IS USED.

FICAM PLUS[®]

SYNERGIZED PYRETHRINS

FOR USE ONLY BY SERVICE PERSONS. NOT FOR SALE TO OR USE BY HOMEOWNERS. NOT FOR STORAGE IN THE HOME.

ACTIVE INGREDIENTS:	Percent by Weight
Permethrin	28.4%
Pyrethrins	3.0%
Piperonyl Butoxide, Technical	7.6%
Petroleum Distillate	12.8%
INERT INGREDIENTS	47.2%
Total	100.0%

Formulated by U.S. Service, 5752-C
 Registered with EPA, Registration No. 15029-1, 15029-2
 Equipment Use: 15029-1 (EPA/Registration/15029-1) 15029-2 (EPA/Registration/15029-2)

EPA Reg. No. 65939-609

EPA Est. 45539-MI-T

PRECAUTION AT USAGE: Not to be used on lawns, in use on produce, or on other plants and animals. Do not use on plants and animals.

KEEP OUT OF REACH OF CHILDREN. WARNING

STATEMENT OF PRACTICAL TREATMENT

IF SEVERAL INSECT INFESTATIONS OF SEVERAL DIFFERENT SPECIES ARE FOUND IN ONE AREA, THE USER SHOULD APPLY FICAM PLUS TO ALL AREAS OF INFESTATION. FICAM PLUS IS A BROAD SPECTRUM INSECTICIDE WHICH IS EFFECTIVE AGAINST A WIDE RANGE OF INSECTS. IT IS ESPECIALLY SUITABLE FOR THE CONTROL OF COCKROACHES, FLEAS, TICKS, AND SPIDERS. FICAM PLUS DOES NOT DAMAGE PAINTS, PLASTICS, FABRICS OR OTHER SURFACES WHERE WATER OR OTHER LIQUIDS ARE APPLIED.

ANDORAM
 THE NATIONAL INSECT CONTROL BOARD
 3000 W. 12TH AVENUE, DENVER, CO. 80202

IMPORTANT: Read the entire Directions for Use, Precautions and Disclaimer of Warranties before using this product.

GENERAL INFORMATION: FICAM PLUS is a specially formulated wettable powder which gives knockdown and residual control of insect pests infested by it. It is especially suited for control of cockroaches.

FICAM PLUS does not damage paint, plastics, fabrics or other surfaces where water or other liquids are applied.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Thoroughly wash out sprayer and strainer with water and detergent before using FICAM PLUS.

For use by service persons and their immediate surroundings and on modes of transport. Permitted areas of use include, but are not limited to, stores, warehouses, metal buildings, truck stops, schools, nursing homes, hospitals, restaurants, hotels and food manufacturing, processing and servicing establishments and on vessels, aircraft, rail cars, buses, trucks and trailers.

FICAM PLUS is intended for use on insects and mites. Do not use FICAM PLUS on mammals, birds, fish, shellfish, or other animals. Do not use FICAM PLUS on plants, trees, shrubs, or other vegetation. Do not use FICAM PLUS on lawns, lawns, or other areas where it is not intended to be used. Do not use FICAM PLUS on surfaces where it is not intended to be used. Do not use FICAM PLUS on surfaces where it is not intended to be used.

Dosages for Conventional Spray Volumes:

1. One packet of FICAM PLUS mixed in one gallon of water yields 0.002% of FICAM PLUS in the spray.
2. Two packets of FICAM PLUS mixed in one gallon of water yields 0.004% of FICAM PLUS in the spray.

INDOOR USE

Includes both non-resistant and organophosphorus resistant strains, ants, beetles, silverfish and firebrats. Mix one packet of FICAM PLUS per gallon of water and apply as a coarse low pressure spray or coarse fog to the areas, closets, around water pipes, doors and windows, behind and under refrigerators, cabinets, sinks and stoves, the underside of shelves, drawers and shelves and other places where ants enter premises.

Following cleanup applications or where insect infestation is low and where sites are respected and treated at monthly intervals, one packet of FICAM PLUS per two gallons of water.

Complete Area: Mix one packet of FICAM PLUS per gallon of water and apply to all walls, around doors and windows and other places where insects are possible, apply FICAM PLUS to all cracks and crevices.

Tentacles: Mix one packet of FICAM PLUS per gallon of water and apply to voids or channels in damaged wooden members and the foundations, in locations vulnerable to termite attack such as crawl spaces. Drilling may be necessary prior to applying FICAM PLUS to cracks or holes in wood.

Apply FICAM PLUS to all cracks or holes in wood. Do not use FICAM PLUS in place may not result in complete coverage of infested and untreated wood or that in completely occupied areas of structures must be securely plugged.

The purpose of such applications of FICAM PLUS is to kill workers or winged reproductive forms which may be present and to impart a temporary resistance at these places to immediate reinfestation by termites. Such infestation of wood is a violation of Federal law.

Foundation treatment but merely a supplement, particularly where use of other insecticides may pose an odor problem. Do not apply solution until location of nest or air conditioning ducts are known and identified. Extreme caution must be taken to avoid contamination of these structural elements and drywalls.

Cracks and Floor in Buildings: Mix one packet of FICAM PLUS per gallon of water and thoroughly apply spray to infested areas such as pet beds and nesting quarters, nearby cracks and crevices, rugs and floor coverings, and under their edges. Do not use FICAM PLUS on surfaces where it is not intended to be used. Do not use FICAM PLUS on surfaces where it is not intended to be used.

Wasp and Bee:

Mix one packet of FICAM PLUS per gallon of water and apply to all areas where wasps and bees are present. Do not use FICAM PLUS on surfaces where it is not intended to be used. Do not use FICAM PLUS on surfaces where it is not intended to be used.

Spider (Including Black Widow Spider):

Mix one packet of FICAM PLUS per gallon of water and apply to all areas where spiders are present. Do not use FICAM PLUS on surfaces where it is not intended to be used. Do not use FICAM PLUS on surfaces where it is not intended to be used.

Wasp and Bee:

Mix one packet of FICAM PLUS per gallon of water and apply to all areas where wasps and bees are present. Do not use FICAM PLUS on surfaces where it is not intended to be used. Do not use FICAM PLUS on surfaces where it is not intended to be used.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Harmful or fatal if swallowed. May cause eye irritation. May be irritating to skin. Avoid contact with eyes. Skin and clothing, if contact occurs, wash immediately with soap and warm water, and eyes with clean water.

Avoid breathing and working in spray, mist or dust. Avoid contact with eyes, nose, mouth, or clothing. Do not smoke and after spraying or handling equipment, wash hands thoroughly before re-use.

Do not apply to humans, their clothing or bedding. Do not permit children or pets near treated surfaces until with most of completely.

Do not apply to food, feed, or other animal products. Do not apply to food, feed, or other animal products.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Keep out of lawn and garden lawn. Do not contaminate water by cleaning of equipment or disposal. This product is highly toxic to bees exposed to direct treatment or to residues on plants.

STORAGE AND DISPOSAL

Store in a cool, dry place. PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. CONTAINER DISPOSAL: Packet - After insuring that both used packets and all trash containers are empty, crush and dispose of in trash container.

DISCLAIMER OF WARRANTIES: THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, OR OF MERCHANTABILITY, WHICH EXTEND BEYOND THE STATE OF ILLINOIS. THE USER ASSUMES ALL LIABILITY FOR ANY DAMAGE, INCLUDING PERSONAL INJURY, PROPERTY DAMAGE, OR OTHER DAMAGE, WHICH MAY BE CAUSED BY THE USE OF THIS PRODUCT. THE USER ASSUMES ALL LIABILITY FOR ANY DAMAGE, INCLUDING PERSONAL INJURY, PROPERTY DAMAGE, OR OTHER DAMAGE, WHICH MAY BE CAUSED BY THE USE OF THIS PRODUCT.

CAREFULLY AND DO NOT DISCARD UNTIL ALL PRODUCT IS USED.

FICAM[®]

76% WETTABLE POWDER

INSECTICIDE

FOR USE ONLY BY PROFESSIONAL APPLICATORS

ACTIVE INGREDIENT:
 1,3-benzodioxol-4-yl-methylcarbamate) 76%
INERT INGREDIENTS: 24%
 Total 100%

Protected by U.S. Patent No. 3,726,338
 EPA REGISTRATION NO. 45639-1

KEEP OUT REACH OF CHILDREN

WARNING

HARMFUL OR FATAL IF SWALLOWED
 IF SWALLOWED — DRINK 1 OR 2 GLASSES OF WATER IMMEDIATELY. DO NOT INDUCE VOMITING OR GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. GET MEDICAL ATTENTION.
 IF INHALED — KEEP PATIENT STRICTLY AT REST AND TRANSFER TO HOSPITAL IMMEDIATELY TO GET MEDICAL ATTENTION.
 IF ON SKIN — WASH WITH SOAP AND WARM WATER. IF IN EYES — RINSE THOROUGHLY WITH CLEAN WATER.
 NOTE TO PHYSICIAN: CONTAINS ANTIQUINONE ESTERASE INHIBITORS. IN CASES OF ACUTE OR CHRONIC EXPOSURE TO THIS PESTICIDE, SUCH AS FOLIAGE FEEDING TO LIVESTOCK, ADULT DOSES SHOULD BE GIVEN AND REPEATED AS NECESSARY UNTIL FULLY RECOVERED.

BFC Chemicals, Inc.
 4311 Lancaster Pike, P. O. Box 2867
 Wilmington, DE 19805

DIRECTIONS FOR USE

FICAM W does not damage paints, plastics, fabrics or other surfaces where water alone causes no damage.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
 Thoroughly wash out sprayer and strainer with water and detergent before using FICAM W.

For use by Professional Applicators for residual pest control in and on buildings and structures and their immediate surroundings and on modes of transport. Permitted areas of use include, but are not limited to, stores, warehouses, industrial buildings, houses, apartment buildings, hospitals, greenhouses, laboratories, schools, nursing homes, restaurants, motels and food manufacturing, processing and serving establishments and on vessels, aircraft, rail cars, buses, trucks and trailers.

FICAM W is intended for dilution with water for application with hand or power-operated sprayers. Do not use strainers finer than 100 mesh size. When using hand sprayers, fill with the desired volume of water, add FICAM W, close and shake before use to insure proper suspension of the wettable powder. Shake or re-agitate sprayer before use if spraying is interrupted.

Dosage Chart: For Conventional Spray Volumes

Use the scoop provided for measuring from the poly jar.
 Each level scoopful contains 0.4 oz FICAM W, which equals one packet.

Amount of FICAM W	Amount of Water	Concentration of Spray Active Ingredient (a)
1 packet or 1 scoop (0.4 oz) per	Two gallons of water	0.125%
1 packet or 1 scoop (0.4 oz) per	Gallon of water	0.25%
2 packets or 2 scoops (0.8 oz) per	Gallon of water	0.50%
1 pound per	40 gallons of water	0.25%

spray in areas where these pests hide, such as: bookshelves, storage areas, closets, around water pipes, doors and windows, behind and under refrigerators, cabinets, sinks and stoves, the underside of shelves, drawers and similar areas. For ants, apply to ant trails, around doors and windows and other places where ants enter premises.

Carpenter Ants: Mix one packet or one scoop of FICAM W per gallon of water (0.25% a) and apply to ant trails, around doors and windows, and other places where ants enter premises, where possible, apply FICAM W directly to ant nest or infested wood.

Following cleanup applications or where insect infestation is low and where sites are inspected and treated at monthly intervals or less, these insects can be controlled with sprays of one packet or one scoop of FICAM W per two gallons of water (0.125% a).

Termites: Mix one packet or one scoop of FICAM W per gallon of water (0.25% a) and apply to voids or channels in damaged wooden members of a structure, or to cracks, spaces or bearing joints between wooden members of a structure or between stud members and the foundations, in locations vulnerable to termite attack. Drilling may be necessary prior to applying FICAM W to voids, cracks or joint spaces. In vulnerable locations, such as the ends of joists, rafters, and other structural members, such drilling and treating of a timber joist may not result in complete coverage of infested and vulnerable voids or galleries.

The purpose of such applications of FICAM W is to kill workers or winged reproductive forms which may be present in the treated chambers and spaces at the time of treatment and to impart a temporary resistance at these places to termite attack which may recur. In some part, in the prevention of immediate reinfestation by termites. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but merely a supplemental, particularly where use of other insecticides may pose an odor problem.

FICAM W at 0.25% a) has consistently provided control of all of the above insects for 30 and often for 60 days.

Brown Dog Ticks and Fleas in Buildings: Mix one packet or one scoop of FICAM W per gallon of water (0.25% a) and thoroughly apply spray to infested areas such as pet beds and resting quarters, nearby cracks and crevices, rugs and floor coverings, and under their edges, between and under cushions of upholstered furniture and other areas where the pests may be present. Use a 1/2 to 1 cup of water and furniture. Old bedding or mats should be removed and replaced with clean, fresh bedding after treatment of pet areas. Additionally, ensure that pets are treated with an EPA approved product registered for such use. Do not treat pits or their bedding with this product. Do not permit children or pets near treated surfaces until spray has dried completely.

Japanese Beetles — Ailcraft: Mix one packet or one scoop of FICAM W per gallon of water (0.25% a). Apply to areas where beetles may enter or be found such as carpets, under seats, storage areas, closets, around doors, behind and under galley equipment, in cargo holds and wheel wells.

Soldiers: Mix one packet or one scoop of FICAM W per gallon of water (0.25% a) and apply to infested baseboards, window and door frames, corners, pipes, storage localities, attics and under eaves. Make spot applications to other areas over which these pests may crawl.

Wasps and Bees: Mix one packet or one scoop of FICAM W per gallon of water (0.25% a) and thoroughly spray nest and entrance and surrounding areas where insects alight. It is generally advisable to spray the nests in the evening when the insects are less active and have returned to the nest. For best results, check nest carefully one or two days after spraying to insure complete kill. Then remove and destroy nest to prevent emergence of newly hatched insects.

Sawflies, Pillbugs, Ground Beetles, Earwigs, Millipedes, Centipedes and Scorpions: Mix one packet or one scoop of FICAM W per gallon of water (0.25% a) and apply around perimeter of buildings and other structures where these pests may enter premises. Spray baseboards, storage areas and other locations where these pests are found.

er with staining before use. Replacement of food and utensils, insected foodstuffs should be destroyed.

Carpet Beetle (Black, Furniture and Vardick): Mix one packet or one scoop of FICAM W per gallon of water (0.25% a.i). Apply as spot application on and under edges of floor coverings, under rugs and furniture, and in closets or other localities where these insects are found. Entire rugs may be treated.

Note: Where it is desired to obtain a longer period of control of all of the above pests in areas where access is difficult or where deposits will not be quickly removed by cleaning, apply FICAM W at a rate of two packets or two scoops per gallon of water (0.50% a.i). This rate may also be used to quickly reduce heavy infestations of cockroaches.

Washing Clothes Moths: Mix two packets or two scoops of FICAM W per gallon of water (0.50% a.i) and apply spray as spot application in closets and other storage areas where washing clothes moths are found. This treatment is an adjunct treatment and will not control moth larvae already on the clothes. Do not apply FICAM W to clothing. To prevent recurrence of pest infestations, repeat applications as required.

Interior Plants: Refer to the ornamental pests recommendations in the Outdoor Use section.

Food Handling Establishments — Places other than private residences in which food is held, processed, prepared or served.

Food Areas — Application limited to spot or crack and crevice treatments only. For spot treatments, no individual spot will exceed two (2) square feet.

Includes areas for receiving, storage, packing (tanning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas when food exposed and facility is in operation are also considered a food area.

Prepare a spray solution as directed above and apply in small amounts directly into cracks and crevices using equipment capable of delivering a penetration of insecticide in points between different elements of construction between equipment and floors, openings leading to voids and into hidden areas. In such equipment as logs, pans, cockroaches, and sabblerfish, fibertex, equipment and shelves, etc. Spot treating crevices may also be made to areas where top of the above pests hide or inhabit, such as: baseboards, storage areas, closets, around water pipes, doors and windows, behind and under refrigerators, cabinets, sinks, stoves and other equipment, the underside of shelves, drawers and similar areas.

Make applications using low pressure to avoid possible atomization or splashing of the spray. Avoid contamination of food or food processing surfaces.

Non-Food Areas — FICAM W may be used as a crack and crevice treatment or as a spot application in non-food areas. All areas where insects hide or through which insects may enter, should be treated.

This product is authorized for use in non-food areas of Federally Inspected Meat and Poultry Plants.

OUTDOOR USE

Perimeter Treatment — Residual spray for control of ants, bees, brown dog ticks, centipedes, cockroaches, crickets, sawflies (firebrats), fleas, ground beetles, millipedes, pillbugs, scorpions, silverfish, sowbugs, spiders, and wasps. Mix one packet or one scoop of FICAM W per gallon of water (0.25% a.i) and apply as a residual spray or with a paint brush to surfaces of buildings, porches, screens, window frames, eaves, patios, garages and other areas where these pests congregate. To help prevent infestations of buildings by the above pests, treat a band of soil and vegetation 6 to 10 feet wide around and adjacent to the building. Also, treat the building foundation to a height of 2 to 3 feet where pests are active and may find entrance. Mix one packet or one scoop of FICAM W per gallon of water (0.25% a.i) and apply as a coat, spray to thoroughly and uniformly wet the band area.

ants, crickets, centipedes, cockroaches, crickets, sawflies (firebrats), fleas, ground beetles, millipedes, pillbugs, scorpions, silverfish, sowbugs, spiders, and wasps. Mix one packet or one scoop of FICAM W per gallon of water (0.25% a.i) and apply as a residual spray or with a paint brush to surfaces of buildings, porches, screens, window frames, eaves, patios, garages and other areas where these pests congregate. To help prevent infestations of buildings by the above pests, treat a band of soil and vegetation 6 to 10 feet wide around and adjacent to the building. Also, treat the building foundation to a height of 2 to 3 feet where pests are active and may find entrance. Mix one packet or one scoop of FICAM W per gallon of water (0.25% a.i) and apply as a coat, spray to thoroughly and uniformly wet the band area.

Do not use on Cardoon, since plant injury may occur.

IMPORTANT: FICAM W has demonstrated excellent plant safety, however, all species and varieties have not been tested. When treating large numbers of plants of a single variety, treat a few plants and observe prior to full-scale application.

Apply FICAM W to paint of run-off using recommended rates indicated below.

DOSAGES AND DIRECTIONS FOR USE ON ORNAMENTAL PLANTS

PEST

Chrysanthemum aphid, Eastern tent caterpillar, Gypsy moth, Hawthorn scale, Hawthorn scale, Florida wax scale

Brown soft scale, Spruce aphid, Yew scale, Citrus scale
Japanese beetle, Turkey Azalea lecanium, Azalea caterpillar, Texas mealybug (larva), Elm leaf beetle, Willow leaf beetle (larva), Fragile tent maker, Yellow-necked caterpillar, Spring cankerworm, Bagworm, Sycamore leafminer, Minnesota webworm, Osageur root weevil (adults)

Greenhouse whitefly, Citrus thrips (larva), Black vine weevil, Pine aphid, Nandivora pine fly, mealybug
Texas pine weevil, Northern pine weevil, Azalea leafminer, Osageur root weevil

When measuring small amounts of product, use the 1 oz. scoop provided. Spray at the time of moth flight for each generation. Consult local Cooperative Extension authorities regarding time of moth flights in your area.

Ornamental Turf Pests: Apply FICAM W as a coarse, low-pressure spray in sufficient water to obtain complete and uniform coverage of pest-infested areas, according to the directions in the table below.

PEST	DOSAGE	APPLICATION PROCEDURE
White grub or Black turf grass Acanthia	1/2 oz. per 1000 sq. ft.	Apply when pests become active. Use a watering can or sprayer when they are still small; thereafter, use a lawn mower or roller for application. Apply in the evening to prevent evaporation. Apply in the morning to prevent evaporation.
European chaffer	1/2 oz. per 1000 sq. ft.	Apply when pests become active. Use a watering can or sprayer when they are still small; thereafter, use a lawn mower or roller for application. Apply in the evening to prevent evaporation. Apply in the morning to prevent evaporation.
Annual white grub	1/2 oz. per 1000 sq. ft.	Apply when pests are evident. Use a watering can or sprayer when they are still small; thereafter, use a lawn mower or roller for application. Apply in the evening to prevent evaporation. Apply in the morning to prevent evaporation.

Fire Ants: Mix FICAM W as recommended in the table below. Two gallons of mixture are recommended for mounds 6 inches or larger in diameter. Less mixture can be used on smaller mounds; however, thorough penetration of insect galleries must be achieved. Apply a small amount of material around the perimeter of the mound. Apply the remaining volume directly on the mound so as to penetrate insect galleries. A second treatment may be necessary if complete control is not obtained.

Do not disturb the mound prior to application. If possible, make applications after a recent rainfall. Applications may not be as effective during hot, dry periods. If possible, treat all colonies in the vicinity. Also, treat colonies which have not yet constructed a mound.

DOSEAGE TABLE	AMOUNT OF FICAM W	AMOUNT OF WATER
1/2 oz. per 1000 sq. ft.	1 packet	2 gallons
1/4 oz. per 1000 sq. ft.	1/2 packet	2 gallons

WARNING USE PRECAUTIONS

Harmful or fatal if swallowed. May be absorbed through the skin. Avoid contact with eyes, skin and clothing. If contact occurs, wash skin and clothing immediately with soap and warm water, and eyes with clean water.

Avoid breathing and working in spray mist or dust. Wash hands and exposed skin before eating, drinking or smoking and after spraying or handling. Wash all contaminated clothing thoroughly before re-use.

Do not apply to humans, their clothing or bedding. Do not permit children or pets near treated surfaces until spray has dried completely. With proper spraying, FICAM W does not result in a visible deposit. However, if unevenly applied, or sprayed to run-off, a visible deposit may occur on some dark surfaces.

Do not spray where electrical short circuits might result. Do not use on or contaminate fruit, vegetables or other feed or food crops. Do not store near feed or food products and do not contaminate these or food preparation surfaces, dishes, kitchen utensils and food containers with either concentrate or spray.

ENVIRONMENTAL HAZARDS
This product is toxic to wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water. Keep out of lawn and garden. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or to residues on plants.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not re-use empty containers.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse that cannot be used or chemically reprocessed, should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL: Packet — After insuring that both used packets and outer containers are empty, crush and dispose of in trash container. Poly Jar — After insuring that container is empty, triple rinse and dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.

GENERAL: Consult federal, state or local authorities for approved alternative procedures such as limited open burning.

DISCLAIMER OF WARRANTIES: THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE ON OTHER THAN THE STATEMENTS MADE ON THIS LABEL. No agent of BFC Chemicals, Inc. is authorized to make any warranties beyond those contained herein or modify the warranties contained herein. UNDER NO CIRCUMSTANCES SHALL BFC CHEMICALS, INC. BE LIABLE FOR ANY INCIDENTAL OR CONSEQUENTIAL DAMAGES. RESULTING FROM FAILURE TO USE THIS PRODUCT IN ACCORDANCE WITH LABEL DIRECTIONS.