LONG LASTING POWDER

— Use In --

Restaurants Hones

Warehouses Schools Sincery Stores

Hotels Mobile Homes Offices

Street

Hospitals

KEEP OUT OF REACH OF

CHILDREN

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONS

CAUTION

Ships Nursing Homes

1985 ng Kennels

rachis

New Building 2007 Sewers Construction

> Kills Cockroaches, Ants, Silverfish, Palmettobugs and Waterbugs.

ORTHOBORIC ACID.....99% with BORIC ACID

ACTIVE INGREDIENT:

ODORLESS

BORID is a long-lasting powder which is effective as long as it is in place and kept gry BORID is a non-flammable, stainless infestations in homes and other buildings odorless powder that will control cockreach

(Product is filled by weight and may settle in shipping)

HAZARDS TO HUMANS AND DOMESTIC ANIMALS PRECAUTIONARY STATEMENTS

CAUTION

HARMFUL IF SWALLOWED, MAY CAUSE EYE IFRITATION, AVOID EYE CONTACTION.

STATEMENT OF PRACTICAL TREATMENT (#2) Immediately contact Physician or Poison Control Center if swaltowed. If eye contact occurs, flight eyes with plenty of water. Get medical attention if irritation persists.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DOMESTIC APPLICATION

No powder should be visible in living areas after application. Any powder visible after application must be brushed into cracks and crevices or removest. Apply only in areas inaccessible to children and pels. Avoid contamination of feed and foodstuffs. Do not use in serving areas when food is expossed. Do not use in edible product areas. Apply approximately 1 pound for every 1000 squara sel of Irving space.

Apply BORID liberally in areas where cockroaches hide or nest:

Under and behind appliances.
In plumbing voids accessible through holes, around plumbing and drains, in politicities under kitchen and bathroom cabinets, (Note: Many cabinets have a gap at the top of the kitchplate which offers easy access. If not, drill a small hole at the top of the kitchplate) in drawer wetls of cabinets.

in cracks around windows, doors, baseplates, flutures and cabinets. In utility rooms under and behind hot water heaters, washers and dryers. Be sure to dust

plumbing voids.

in attics - around the pennteler and vent pipes, above kitchens and baths. Around the perimeter of dropped ceilings, into pits of elevator shalfs, durithwallers and tresh chutes. Under and behind pallets and base in storerooms.

IET WEIGHT

DIRECTIONS FOR USE

WITHIN FOOD AREAS AND NON-FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS.)

FOOD HANDLING ESTABLISHMENTS: Places other than private residences in which exposed tood is held, processed, prepared or served.

FOOD AREAS APPLICATION IS LIMITED TO CRACK AND CREVICE TREATMENT ONLY. Apply in small amounts directly into cracks and crevices using a bellows duster or other suitable equipment capable of applying insecticide directly. Includes areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, daties, edible oits, syrups). Serving areas, when food is exposed and facility is in operation, also would be considered a food area.

NON-FOOD AREAS: Includes garbage rooms, lavatories, floor drains (to sewer entries and restitutes), offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage rather canning or bottling). Apply to baseboard areas, around water pipes, surfaces behind and beneath sinks, lockers, tables, pallets and similar areas where insects hide or through which they

Care should be taken to avoid depositing the product onto surfaces which are exposed or introducting the material into the air. Avoid contamination of food or food processing surfaces. APPLICATIONS OF THIS PRODUCT IN THE FOOD AREA OF FOOD HANDLING ESTABLISHMENTS.

THER THAN AS A CRACK AND CREVICE TREATMENT ARE NOT PERMITTED.

(S)SEWER TREATMENT: Apply this product at a rate not to exceed 8 ibs per 3000 sq ft using portable blowers or air pressure tanks.

PRETREATMENT OF NEW CONSTRUCTION: Apply powder to base plates and plumbing voids, at the rate of 4 lbs per 1000 sq ft. In artics apply powder around perimeter of affics and around vent stacks at the rate of 4 lbs. per 1000 sq. ft.

CARPENTER ANTS: Locate source of infestation. Using power duster, inject powder for 3-5 secunds into insect tunnels and cavilles.

STORAGE & DISPOSAL

On not contaminate water, food or feed by storage or disposal. STORAGE: Store in original container away from food and feeds utfa. Keep products in a locked storage area.

DISPOSAL: Do not reuse this container. Wrap and place in trash collector.

EPA Reg. No. 9444-129

WATERBURY COMPANIES, INC.

EPA Est. No. 9444-LA-1